



Belgian Cancer Registry

CANCER FACT SHEET 2022

HEAD & NECK CANCER

ICD-10 C00-C14; C30-C32



Key facts





- 5th most common cancer in males
- 9th most common cancer in females
- 2,721 new diagnoses in 2022
- 679 deaths due to H&N cancer in 2021
- 5-year net survival of **58.0%**




Lifetime risk (0-84 years)



3.2 in 100 males

1.2 in 100 females






- Median age at diagnosis for H&N cancer is **66 years**
- Known risk factors include smoking, alcohol consumption and human papilloma virus (HPV) infection
- There is a **higher risk** of H&N cancer diagnosis **closer to the French border**
- **More than half** of H&N cancers are located in the oral cavity or oropharynx


Incidence by sub-location, N(%)	Males		Females	
Lip	21	(1.1%)	16	(2.1%)
Oral cavity	466	(24.0%)	260	(33.4%)
Oropharynx	540	(27.8%)	214	(27.5%)
Nasopharynx	42	(2.2%)	27	(3.5%)
Hypopharynx	222	(11.4%)	54	(6.9%)
Larynx	456	(23.5%)	94	(12.1%)
Nasal cavity and paranasal sinuses	96	(4.9%)	44	(5.7%)
Salivary glands	87	(4.5%)	69	(8.9%)
Other and ill-defined sites	13	(0.7%)	0	(0.0%)



Stage distribution by main sub-location



HPV status in oropharyngeal cancer




Squamous cell carcinomas of the oropharynx


- Stage distribution is dependent on **sub-location**
- At least **one in three** oropharyngeal cancers is related to HPV infection
- Prognosis is dependent on HPV status, available on page 5
- More information on the anatomy of H&N is available on page 2




Age-standardised incidence and mortality, 2004-2022





- H&N cancer is **more common in males** than in females; male to female ratio of 2.5
- Risk of a H&N cancer diagnosis **in males is decreasing** with an average annual percentage change of **-1.1%**
- Risk of a H&N cancer diagnosis **in females is increasing** with an average annual percentage change of **+1.0%**



Incidence by main sub-location, 2013-2022






- **5-year net survival has been improving** for males and females in the last 15 years
- Diagnosis in **an early stage** is associated with a **better prognosis**
- More than **17,000 people** are living with the consequences of H&N cancer

Additional detailed information (including prevalence) can be found in the appendix of the Cancer Fact Sheet and on the website of the Belgian Cancer Registry.



5-year net survival, 2018-2022, % (95% CI)		
Lip	85.6%	(68.5%; 93.8%)
Oral cavity	56.3%	(53.8%; 58.8%)
Oropharynx	53.2%	(51.0%; 55.4%)
Nasopharynx	72.7%	(65.0%; 79.0%)
Hypopharynx	34.9%	(31.4%; 38.4%)
Larynx	67.6%	(64.4%; 70.6%)
Nasal cavity and paranasal sinuses	63.4%	(58.0%; 68.4%)
Salivary glands	77.6%	(67.5%; 84.8%)





- **Absolute numbers (N):** The number of newly registered cancer diagnoses observed for a given period of time. All figures and numbers in this cancer fact sheet are based on diagnoses of Belgian residents.
- **Crude Rate (CR):** The crude rate is obtained by dividing the absolute number of diagnoses (N) by the corresponding population size at risk (N/100,000).
- **ESR2013:** Incidence rates standardised to the 2013 revised European Standardised Population (ESP): standardisation is needed to accommodate for differences in population size and age distribution (over time or among regions). An important factor in interpreting trends in cancer incidence is population ageing, as cancer is an age-dependent disease. For a higher proportion of elderly people in the population, a higher total number of cancer diagnoses can be expected for the same cancer risk. When only absolute numbers (N) or Crude Rate (CR) results are used, a misleading picture of the actual changes in the risk of a cancer diagnosis could be obtained. Therefore, direct standardisation is needed to evaluate the evolution of the risk of cancer diagnosis over time or among regions.
- **Cancer maps:** Cities with at least 150,000 inhabitants are directly represented on the map as circles with a diameter relative to the population size, and a colour shading indicating the actual calculated ESR2013 in that city. The 19 municipalities of the Brussels Capital Region (more than 1,000,000 inhabitants) are divided in three separate circles, based on socio-economic parameters. The socio-economic status is lowest in the westernmost circle and highest in the easternmost circle. Methodological information is available in 'Cancer burden in Belgium 2004-2017, Belgian Cancer Registry, Brussels, 2020'.
- **Stage:** Cancers are reported with a stage, labelled with a Roman numeral with IV being the most advanced stage. The stage is based on the T-category (extent of the tumour), the N-category (absence or presence and extent of the regional lymph node metastasis) and the M-category (absence or presence of distant metastasis). Stage is reported as a combination of both clinical and pathological stage with priority given to the pathological stage. Clinical information about distant metastases (cM) will always be taken into account, and in case of neo-adjuvant therapy, priority is given to the clinical stage. If stage is unknown, not applicable or not submitted to the Belgian Cancer Registry, the stage is reported as 'unregistered stage'. Stage is reported according to the TNM 8th edition: J.D. Brierley, M.K. Gospodarowicz, Ch. Wittekind. TNM Classification of Malignant Tumours, 8th edition: UICC, 2017.
- **Net survival:** often also called the relative survival, is an estimate of the survival probability when other causes of death beside the cancer type(s) under study are excluded. As examples of other causes of death, patients with the cancer type(s) under study could also die because of an accident or unrelated cardiac conditions, etc.
- **95% CI:** 95% confidence intervals are indicated with a shaded band or whiskers in the figures. The 95% CI is a range of values that has 95% chance to contain the true mean value.
- **HPV:** Human Papillomavirus is a sexually transmitted infection and a risk factor for the development of some H&N cancers.
- **H&N cancer is divided in sub-locations according to ICD-10 codes:**
 - **Lip:** C00
 - **Oral cavity:** C02-C05.0; C06
 - **Oropharynx:** C01; C05.1-C05.9; C09-C10
 - **Nasopharynx:** C11
 - **Hypopharynx:** C12-C13
 - **Larynx:** C32
 - **Nasal cavity and paranasal sinuses:** C30-C31
 - **Salivary glands:** C07-C08
 - **Other and ill-defined sites:** C14

**Mortality statistics in Belgium are collected and managed by the three Regions (Flemish Region: Agentschap Zorg en Gezondheid; Brussels-Capital Region: Observatorium voor Gezondheid en Welzijn van Brussel-Hoofdstad/ l'Observatoire de la Santé et du Social de Bruxelles-Capitale; Walloon Region: Agence Wallonne de la Santé, de la Protection sociale, du Handicap et des Familles (AVIQ). The Directorate General Statistics Belgium is responsible for collecting and merging the data coming from the regional agencies. Mortality data used in this cancer fact sheet are collected from the Directorate General Statistics Belgium and encompasses the period 2004-2021).*

Recommended reference: Cancer Fact Sheets 2022, Belgian Cancer Registry (BCR), 2024